

EXPLORING THE CHARACTERISTICS OF DUPLICATION AND SWITCHING IN DIVISOR CORDIAL GRAPHS

¹Dr. John Smith and ²Dr. Emily Johnson

¹Department of Mathematics, North Valley Institute of Technology, Springfield, Illinois, USA

²Department of Mathematics, Greenfield University, Pinehill, New Hampshire, USA

Abstract:

: Cordial graphs, a class of graphs introduced in 1987, have been a topic of interest in graph theory. This paper explores various properties and characteristics of cordial graphs, including their relationship with other types of graphs and specific conditions for their cordiality. The study also investigates cordiality in different graph structures and provides insights into the conditions under which certain graphs can be classified as cordial.

Keywords: Cordial graphs, binary labeling, graph theory, graph properties, combinatorial mathematics.

1. Introduction

By a graph, we mean a finite undirected graph with out loops and multiple edges. For terms not defined here, we refer to Harary [7]. Cordial graph was first introduced by I. Cahit [1] in 1987 as a weaker version of graceful and harmonious graphs and was based on $\{0, 1\}$ binary labeling of vertices. He showed that (i) every tree is cordial (ii) C_n is cordial if and only if $n \leq 3$ (iii) C_n is cordial for all r and s (iv) the wheel W_n is cordial if and only if $n \equiv 3 \pmod{4}$ (v) C_n is cordial if and only if $n \not\equiv 2 \pmod{4}$ (vi) an Eulerian graph is not cordial if its size is congruent to 2 modulo 4. Other types of cordial graphs are considered in [3, 7, 10, 11]. For more related results on cordial graphs, one can refer to Gallian [6].

Definition 1.1 [1]

A binary vertex labeling of graph $G(V, E)$, where each edge uv is labeled with $|f(u) - f(v)| \pmod{2}$, is called a **cordial labeling** if $|f^{-1}(0) - f^{-1}(1)| \leq 1$ and $|e_0 - e_1| \leq 1$, where $f^{-1}(i)$ denote the number of vertices labeled with i under f and e_i denote the number of edges labeled with i , where $i = 0, 1$. A graph G is called cordial if it admits a cordial labeling.

Definition 1.2 [15]

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$ and let $f: V(G) \rightarrow \{0, 1\}$. Define on $V(G)$ by $f^*(i) = \sum_{f^{-1}(j)=i} f(j) \pmod{2}$. The function f is called an **E-cordial labeling** of G if $|f^{-1}(0) - f^{-1}(1)| \leq 1$ and $|f^*(0) - f^*(1)| \leq 1$. A graph is called **E-cordial** if it admits E-cordial labeling.

In 1997 Yilmaz and Cahit [15] have introduced E-cordial labeling as a weaker version of edge-graceful labeling. More results are seen in [15,4]

Definition: 1.3 [4]

A **prime cordial labeling** of a graph G with vertex set V is a bijection f from V to $\{1, 2, 3, \dots, |V|\}$ such that if each edge uv is assigned the label 1 if $\gcd(f(u), f(v)) = 1$ and 0 if $\gcd(f(u), f(v)) > 1$, then the number of

edges having label 0, and the number of edges having label 1, differ by at most 1. Sundaram et. al [9] has introduced the notion of prime cordial labeling and proved some graphs are prime cordial

Definition: 1.4

The **fan** F_n is the graph obtained by taking $(n - 2)$ concurrent chords in cycle C_n . The vertex at which all the chords are concurrent is called the **apex vertex**. More precisely, $C_n + K_1$ is called the **fan** F_n .

Definition: 1.5

A **helm** H_n , $n \geq 3$ is the graph obtained from the wheel W_n by adding a pendant edge at each rim vertex.

Definition: 1.6

The **flower** F_n is the graph obtained from a helm H_n by joining each pendant vertex to apex vertex of the helm.

Definition: 1.7

The **closed helm** CH_n is the graph obtained from a **helm** H_n by joining each pendant vertex to form a cycle.

Definition 1.8[11]

Let $G = (V, E)$ be a simple graph and $f: V \rightarrow \{1, 2, 3, \dots, |V|\}$ be a bijection. For each edge uv , assign the label 1 if either $f(u) \mid f(v)$ or $f(v) \mid f(u)$ and the label 0 otherwise. Then f is called a **divisor cordial labeling**. A graph with a divisor cordial labeling is called

divisor cordial graph.

R.Varatharajan, S.Navaneethakrishnan and K.Nagarajan [13], introduced the concept of divisor cordial and proved the graphs such as path, cycle, wheel, star and some complete bipartite graphs are divisor cordial graphs and in [14], they proved some special classes of graphs such as full binary tree, dragon, corona, $*$, and $*$, are divisor cordial. We proved in [8] that some special graphs such as switching of a vertex of cycle, wheel, helm, duplication of arbitrary vertex of cycle, duplication of arbitrary edge of cycle, split graph of C_n , W_n , H_n , are divisor cordial graphs.

Labeled graph have variety of application in coding theory, particularly for missile guidance codes, design of good radar type codes and convolution codes with optimal auto correlation properties. Labeled graph plays vital role in the study of X-ray crystallography, communication networks and to determine optimal circuit layouts.

In this paper, we prove that fan graph, switching of a pendant vertex of a helm graph, switching of a vertex of flower graph, switching of closed helm graph and also duplication of a arbitrary vertex by an edge of a fan are divisor cordial.

2. Main results

Theorem: 2.1

The fan F_n is a divisor cordial graph.

Proof:

Let v be the apex vertex and $v_1, v_2, \dots, v_{n-2}$ be the other vertices of the path in fan F_n . Then $p = n + 1$ and $q = 2n - 1$. We define vertex labeling $f: V \rightarrow \{1, 2, \dots, p\}$ as follows.


$$f(v) = 1$$

$$f(v_i) = i + 1; 1 \leq i \leq n-2$$

Since 1 divides any integers, the edges receive label 1 and the consecutive numbers does not divide each other, so the $(n - 1)$ edges in path receive label 0.

Now we observe that $f(v) = 1$; $f(v_1) = 2$.

Hence $f(v) - f(v_1) = 1$.



$$\begin{aligned} () &= 2 \\ () &= 2 + 1; \lceil - \rceil \leq \leq \\ () &= 2 + 1; 1 \leq \leq \lfloor - \rfloor \\ () &= 2(+ 1); \lceil - \rceil \leq \leq \end{aligned}$$

As the above case, we observe that $2 - 1 + \lceil - \rceil$ edges receive label 1.

That is, $(1) = \text{---}$ edges receive label 1 and $(0) = \text{---}$ edges receive label 0

Hence, $(0) - (1) = 1$

From both the cases $| (0) - (1) | \leq 1$

Hence switching of a pendant vertex of is a divisor cordial graph. ■




Figure 2: Switching of a pendant vertex in

Note: 3.3

Switching of a vertex of inner cycle of a helm , the graph becomes disconnected.

Theorem: 3.4

The graph obtained by switching of a vertex of flower graph is divisor cordial.

Proof:

Let w be the apex vertex; be the vertices of cycle and ... be the pendant vertices. We define $f: V(G) \rightarrow \{1, 2, \dots, p\}$

Case (i) Switching of a vertex , $\leq \leq$

Without loss of generality let us assume that the rim vertex is switched. Then $p = 2n + 1$, $q = 3(2n + 1)$

$$\begin{aligned} () &= 1 \\ () &= 2 \\ () &= 2(+ 1); 1 \leq \leq - 1 \\ () &= 2 + 1; 1 \leq \leq - 1 \\ () &= () + 2 \end{aligned}$$

Since v is given label 1 the edges adjacent to v receives label 1 the apex vertex is labeled with 2, and $1 \leq i \leq n$ is labeled with even integers, since they divide each other, that edges receive label 1. Hence $(2n - 1 + n)$ edges label 1 and other $(3n - 1)$ edges receive label 0.

That is, $(1) = (0) = (3n - 1)$

Therefore, $(0) - (1) = 0$




Figure 3: Switching of a pendant vertex of $F_{n/2}$

Case (ii) Switching of a vertex $v, 1 \leq v \leq n$

Let us assume that the vertex v is switched. Here $p = 2n + 1$ and $q = 6n - 8$.

$$(v) = 1$$

$$(v) = 2; 1 \leq v \leq n$$

$$(v) = 2 + 1; 1 \leq v \leq n$$

Since the apex vertex is given label 1, the $(2n - 1)$ edges incident to it receives label 1. The switched vertex is given label 2, and so the $(n - 3)$ vertices incident to v also receives label 1. Hence $(n - 3) + (2n - 1) = 3n - 4$ edges receives label 1 and other $(3n - 4)$ edges receives label 0

That is, $(1) = (0) = (3n - 4)$

Therefore, $|(0) - (1)| = 0$

From both the cases $|(0) - (1)| \leq 1$. Hence graph obtained by switching of a vertex of flower graph is divisor cordial.




Figure: 4: switching of a vertex in

Case (iii) Switching of the apex vertex w , the graph becomes disconnected. ■

Theorem: 3.5

The graph obtained by switching of a vertex of a closed helm graph is integer cordial.

Proof:

Let w be the apex vertex; $v_1, v_2, \dots, v_{2n-1}$ be the vertices of inner cycle and $u_1, u_2, \dots, u_{2n-1}$ be the vertices of outer cycle. Define $f: V \rightarrow \{1, 2, \dots, p\}$

Case (i) Switching of a vertex v_i ; $1 \leq i \leq 2n-1$

Here $p = 2n + 1$ and $q = 6(n - 1)$. Without loss of generality let us assume the vertex v_1 is switched. The labeling is as follows:

$$f(v_1) = 2; f(v_i) = 2i - 1$$

$$f(u_i) = 2(i + 1); 1 \leq i \leq 2n - 1$$

$$f(w) = 2 - 1; 1 \leq i \leq 2n - 1$$

Since the vertex v_1 is given label 1, the $(2n - 3)$ edges incident to it receive label 1. Similarly the apex vertex w is given label 2 and the vertices of inner cycle v_i are given even integers and so $(n - 2)$ edges and the edge (v_1, w) receives label 1. Other edges receive label 0.

That is, $e(1) = e(0) = 3(n - 1)$




Figure: 5 Switching of w in C_{19}

Case (ii) Switching of a vertex w ; $1 \leq w \leq 19$

Without loss of generality let us assume the vertex w is switched. Here $p = 2n + 1$ and $q = 6n - 8$

The labeling is as follows:

$$(1) = 2; (2) = 1; (3) = 2 + 1$$

$$(4) = 2; 2 \leq i \leq 19$$

$$(i) = 2 - 1; 2 \leq i \leq 19. \text{ Now interchange } 1 \text{ and } 2.$$

From the above labeling, we observe that $(3n - 4)$ edges receive label 1 and $(3n - 4)$ edges receive label 0.

That is, $(1) = (0) = 3 - 4$.




Figure: 6 Switching of vertex w in C_{19}

Case (iii) Switching of apex vertex w

Let the apex vertex w be switched. Then $p = 2n + 1$ and $q = 4n$.

Subcase (i) $n \not\equiv 2 \pmod{4}$

The labeling is as follows:

$$(1) = 1$$

The vertices of inner cycle $v_1, v_2, \dots, v_{19}$ and rim vertices $u_1, u_2, \dots, u_{19}$ be labeled in the following order

$$2, 2 \times 2, 2 \times 2, \dots, 2 \times 2$$

$$3, 3 \times 2, 3 \times 2, \dots, 3 \times 2 \quad \dots \dots \dots (2)$$

$$5, 5 \times 2, 5 \times 2, \dots, 5 \times 2$$

$$\dots \dots \dots$$

Where $(2 - 1) \times 2 \leq$ and $\geq 1, > 0$. We observe that $(2 - 1) \times 2$ divide $(2 - 1) \times 2; (<$

) and $(2 - 1) 2$ does not divide $(2m+1)$. Now interchange the labels of and .
The remaining pendant vertices are labeled continuously other than the above labels.
From the labeling, we observe that $2n$ edges receives label 1 and $2n$ edges receive label 0.
That is, $(1) = (0) = 2$.

Subcase (ii) $n \equiv 2 \pmod{4}$

The labeling is as follows:

$$f(w)=1$$

The vertices of inner cycle , , ... and pendant vertices , , ... be labeled in the following order

$$2, 2 \times 2, 2 \times 2, \dots 2 \times 2$$

$$3, 3 \times 2, 3 \times 2, \dots 3 \times 2 \dots \dots \dots (2)$$

$$5, 5 \times 2, 5 \times 2, \dots 5 \times 2 \dots \dots \dots$$

Where $(2 - 1) 2 \leq$ and $\geq 1, > 0$. We observe that $(2 - 1) 2$ divide $(2 - 1) 2$; ($<$) and $(2 - 1) 2$ does not divide $(2m+1)$.

The remaining pendant vertices are labeled continuously other than the above labels.

From the labeling we observe that $2n$ edges receives label 1 and $2n$ edges receive label 0.

That is, $(1) = (0) = 2$.

From all the cases $| (0) - (1) | = 0$

Hence the graph obtained by switching a vertex is divisor cordial. ■




Figure: 6 Switching of apex vertex in

4. Duplication of a vertex and duplication of an edge.

Definition: 4.1 [12]

Duplication of a vertex by a new edge = in a graph G produces a new graph such that
 $(') = \{ , '' \}$ and $('') = \{ , ' \}$

Definition: 4.2[12]

Duplication of an edge = by a vertex in a graph G produces a new graph such that
 $(') = \{ , \}$.

Theorem: 4.3

Duplication of a vertex by an edge in a fan graph is divisor cordial graph.

Proof:

Let v be the apex vertex and $v_1, v_2, \dots, v_n$ be the other vertices of fan and let w and u be the newly added vertex. Then $p = n + 3$ and $q = 2(n + 1)$. We define vertex labeling as $f: V \rightarrow \{1, 2, \dots, p\}$ as follows.

Case (i)

Let the apex vertex v be duplicated.

Subcase (i) n is even

$$f(v) = 2$$

Let $v_1, v_2, \dots, v_n$ be the vertices of path P_n . We label these vertices as follows:

$$\begin{aligned} &1, 2 \times 2, 2 \times 2, \dots, 2 \times 2 \\ &3, 3 \times 2, 3 \times 2, \dots, 3 \times 2 \quad \dots \dots \dots (2) \\ &5, 5 \times 2, 5 \times 2, \dots, 5 \times 2 \end{aligned}$$

where $(2 - 1)^2 \leq$ and $\geq 1, > 0$. We observe that $(2 - 1)^2$ divide $(2 - 1)^2$; ($<$) and $(2 - 1)^2$ does not divide $(2m+1)$. The remaining vertices are given other labels up to p . Then $(n + 1)$ edges receives label 1 and $(n + 1)$ edges receives label 2.

$$\text{That is, } f(1) = f(0) = (n + 1)$$

$$\text{Therefore, } |f(0) - f(1)| = 0$$

Subcase (ii) n is odd

We label as above case. The vertices of path is labeled as

$$\begin{aligned} &1, 2 \times 2, 2 \times 2, \dots, 2 \times 2 \\ &3, 3 \times 2, 3 \times 2, \dots, 3 \times 2 \quad \dots \dots \dots (2) \quad 5, 5 \times 2, 5 \times 2, \dots, 5 \times 2 \\ &\dots \dots \dots \end{aligned}$$

where $(2 - 1)^2 \leq$ and $\geq 1, > 0$. We observe that $(2 - 1)^2$ divide $(2 - 1)^2$; ($<$) and $(2 - 1)^2$ does not divide $(2m+1)$.

$$\text{Then } f(1) = f(0) = (n + 1)$$

$$\text{Therefore, } f(0) - f(1) = 0$$

Case(ii)

Let any of the vertex v_i ; $1 \leq i \leq n$ be duplicated.

Subcase (i) n is even

$f(v) = 2$; $f(v_i) =$ and $f(u) =$; where and are prime numbers Let, $v_1, \dots, v_n$ be the vertices of path P_n . We label these vertices as follows:

$1, 2 \times 2, 2 \times 2, \dots, 2 \times 2$
 $3, 3 \times 2, 3 \times 2, \dots, 3 \times 2$ (2)
 $5, 5 \times 2, 5 \times 2, \dots, 5 \times 2$

Where $(2 - 1)2 \leq$ and ≥ 1 , > 0 . We observe that $(2 - 1)2$ divide $(2 - 1)2$; ($<$) and $(2 - 1)2$ does not divide $(2 + 1)$. Remaining vertices are given other labels other than and.

Subcase (ii) n is odd

We label as in the above case. The vertices of path are labeled upto $\lfloor \frac{n}{2} \rfloor, \lfloor \frac{n}{2} \rfloor \times 2, \lfloor \frac{n}{2} \rfloor \times 2, \dots, \lfloor \frac{n}{2} \rfloor \times 2$

From the above cases, we observe that $f(1) = f(0) = (n + 1)$

Therefore, $f(0) - f(1) = 0$




Figure: 6 Duplication of the vertex in

Case (iii) If the vertex is duplicated then the labeling as follows;

$f(n) = 2$; $f(n) =$; $f(n) = 1$; " = Where and are prime numbers.

Other labels are given as in above cases. From the labeling, we observe that $f(1) = f(0) = (n + 1)$

Therefore, $f(0) - f(1) = 0$

From all cases $f(0) - f(1) \leq 1$.

Hence duplication of a vertex by an edge is a divisor cordial graph. ■

References

I. Cahit, Cordial graphs, A weaker version of graceful and harmonious graphs, ArsCombinatoria, 23, 1987, pp 201208.

I. Cahit, Recent results, and Open Problems on Cordial graphs, Contemporary methods in Graph. Theory, R.

Bodendrek.(Ed.) Wissenschaftsverlog, Mannheim,1990, pp. 209-230. I.

Cahit , H- Cordial Graphs, Bull. Inst. Combin.Appl; 18 (1996) 87-101.

Devaraj J, On Edge-Cordial Graphs. Graph Theory Notes of New York, XLVII, (2004), 14-18.

J.A.Gallian, A dynamic survey of graph labeling, Electron.J.Combin.5 (2000) 1-79.

F.Harary, Graph Theory, Addison- Wesly, Reading Mass 1972.

M.Hovey, A-Cordial Graphs, Discrete Math.1991 (3), pp 183-194

P.Maya and T. Nicholas, Some new families of divisor cordial graph, Annals of Pure and Applied. Mathematics, Vol 5, No. 2, 2014,125-134

M.Sundaram, R.PonrajandS. SomuSundaram, Prime Cordial labeling of graphs, journal of Indian Academy of Mathematics, 27 (2005) 373 – 393 A.Unveren and I.Cahit, M-Cordial Graphs, Pre-print

S.K.Vaidya and LekaBijukumar, Some New families of E-cordial graphs, Journal of Mathematics Research, Vol. 3, No. 4, November 2011.

S.K.Vaidya and N.B.Vyas, E-cordial labeling in the context of switching of a vertex International Journal Of Advanced Computer and mathematical Forum, 4 (31) (2009) 1543-1553.

R.Varatharajan, S.Navaneetha Krishnan, K.Nagarajan, Divisor cordial graphs, International J. Math.Combin.Vol 4 (2011)15 – 25.

R.Varatharajan, S.Navaneetha Krishnan, K.Nagarajan, Special Classes of Divisor cordial, Graphs, International Mathematical Forum Vol 7, (2012), no. 35, 1737 – 1749

R.Yilmaz and I.Cahit, “E-Cordial graphs”, ArsCombin., Vol 46, pp.251 – 266 , 1997 [5]